



US006110696A

United States Patent [19]

Brown et al.

[11] Patent Number: 6,110,696

[45] Date of Patent: *Aug. 29, 2000

[54] ELECTROCHEMICAL ENZYME ASSAY

- [75] Inventors: **Mary E. Brown**, Indianapolis, Ind.; **Hans-Joachim Guder**, Grunstadt, Germany; **John G. R. Hurrell**, Carmel, Ind.; **Lance S. Kuhn**, Fishers, Ind.; **Robert J. McEnroe**, Noblesville, Ind.; **Rebecca W. Muddiman**, Indianapolis, Ind.; **Mary Luann Ochs**, Fishers, Ind.
- [73] Assignees: **Roche Diagnostics Corporation**, Indianapolis, Ind.; **Boehringer Mannheim GmbH**, Germany

[*] Notice: This patent issued on a continued prosecution application filed under 37 CFR 1.53(d), and is subject to the twenty year patent term provisions of 35 U.S.C. 154(a)(2).

This patent is subject to a terminal disclaimer.

[21] Appl. No.: 08/494,668

[22] Filed: Jun. 26, 1995

Related U.S. Application Data

- [63] Continuation-in-part of application No. 08/113,548, Feb. 27, 1993, Pat. No. 5,427,912.
- [51] Int. Cl.⁷ G01N 33/535
- [52] U.S. Cl. 435/7.6; 204/403; 435/18; 435/23; 435/24; 435/287.2; 435/287.7; 435/287.9; 435/817
- [58] Field of Search 435/7.6, 18, 23, 435/24, 817, 287.2, 287.7, 287.9; 204/403

[56] References Cited

U.S. PATENT DOCUMENTS

- 4,378,428 3/1983 Farina et al. 435/7
4,708,929 11/1987 Henderson 435/7
4,785,080 11/1988 Farina et al. 530/402
4,830,959 5/1989 McNeil et al. 435/7
4,912,041 3/1990 Batchelor et al. 435/101
4,950,612 8/1990 Khanna et al. 436/505
4,956,274 9/1990 Khanna et al. 435/7
4,963,245 10/1990 Weetall 204/403
5,032,503 7/1991 Khanna et al. 435/7.6
5,037,735 8/1991 Khanna et al. 435/7.6
5,106,950 4/1992 Farina et al. 530/345
5,120,653 6/1992 Henderson 435/252.33
5,124,253 6/1992 Foulds et al. 435/21
5,202,233 4/1993 Herrmann et al. 435/7.4
5,212,081 5/1993 Coty et al. 435/188
5,223,393 6/1993 Khanna et al. 435/7.6
5,229,282 7/1993 Yoshioka et al. 435/177
5,244,785 9/1993 Loor et al. 435/7.6
5,427,912 6/1995 Brown et al. 435/7.6

5,444,161 8/1995 Manning et al. 536/4.1

FOREIGN PATENT DOCUMENTS

- 0 125 139 A2 11/1984 European Pat. Off. G01N 33/54
0 150 999 A2 8/1985 European Pat. Off. G01N 33/53
0 328 380 A2 8/1989 European Pat. Off. G01N 27/00
0350808 1/1990 European Pat. Off. .
0 546 536 A1 6/1993 European Pat. Off. C12Q 1/00
86/03837 7/1986 WIPO G01N 33/53
86/04926 8/1986 WIPO C12Q 1/00
90/13569 11/1990 WIPO C07K 7/10
91/16630 10/1991 WIPO G01N 33/531
9506115 3/1995 WIPO .

OTHER PUBLICATIONS

Y. Xu et al., Heterogeneous Enzyme Immunoassay of Alpha-Fetoprotein in Maternal Serum by Flow-Injection Amperometric Detection of 4-Aminophenol, (1990) 1941–1944, Clin. Chem. 36/11.

Thompson et al., Comparison of Methods for Following Alkaline Phosphatase Catalysis: Spectrophotometric versus Amperometric Detection, (1991) 90–95, Analytical Biochemistry 192.

Gil et al., Competitive Heterogeneous Enzyme Immunoassay for Theophylline by Flow-Injection Analysis with Electrochemical Detection of p-Aminophenol, (1990) 662–665, Clin. Chem. 36/4.

Jenkins et al., The Use of Ion Pairing Reagents to Reduce Non-specific Adsorption in a Solid Phase Electrochemical Enzyme Immunoassay, (1990) 99–104, Contributed Article vol. 13 No. 2.

Wright et al., Sequestration Electrochemistry: The Interaction of Chlorpromazine and Human Orosomucoid, (1988) 290–293 Analytical Biochemistry 171.

Jenkins et al., Extending the Detection Limit of Solid-Phase Electrochemical Enzyme Immunoassay to the Attomole Level, (1988) 292–299 Analytical Biochemistry 168.

Primary Examiner—David Saunders

Attorney, Agent, or Firm—Roche Diagnostics Corporation

[57]

ABSTRACT

A diagnostic kit, method, and apparatus for electrochemically determining the presence or concentration of an analyte in a sample. A mixture is formed which includes the sample, an enzyme acceptor polypeptide, an enzyme donor polypeptide, and a labeled substrate. The enzyme donor polypeptide is capable of combining with the enzyme acceptor polypeptide to form an active enzyme complex. The formation of such the active enzyme complex is responsive to the presence or concentration of the analyte in the fluid sample. The active enzyme hydrolyzes the labeled substrate, resulting in the generation of an electroactive label, which can then be oxidized at the surface of an electrode. A current resulting from the oxidation of the electroactive compound can be measured and correlated to the concentration of the analyte in the sample.

31 Claims, 5 Drawing Sheets